

# Future change of western North Pacific Typhoons: Projections by a 20-km-mesh global atmospheric model

Hiroyuki Murakami (JAMSTEC/MRI),  
Bin Wang (University of Hawaii/IPRC),  
and Akio Kitoh (MRI)

Reference:


Murakami, H., B. Wang, and A. Kitoh: Future change of western North Pacific typhoons: Projections by a 20-km-mesh global atmospheric model. *J. Climate*, 24, 1154-1169.

## Review and Motivations

- Among many studies investigating future changes in tropical cyclone (TC) activities, few investigated TC track changes.
- Bengtsson et al. (2007, *Tellus*) suggests that future changes in TC density are different between T63 and T213, indicating high-resolution model is necessary for accurate future projections.
- The present study examines future changes in TC tracks, especially in the western North Pacific (WNP), simulated by the 20-km-mesh MRI/JMA-AGCM.


# Model Specifications

	MRI-AGCM 3.1 (Mizuta et al. 2006)
Horizontal resolution	TL959 (20km)
Vertical resolution	60 levels (top at 0.1hPa)
Time integration	Semi-Lagrangian
Time step	6minutes
Cumulus convection	Prognostic Arakara-Schubert
Cloud	Smith (1990)
Radiation	Shibata and Aoki (1989) Shibata and Uchiyama(1992)
GWD	Iwasaki et al. (1989)
Land surface	SiB ver0109(Hirai et al.2007)
Boundary layer	MellorYamada Level2
Aerosol (direct)	Sulfate aerosol
Aerosol (indirect)	No



# Experimental Designs

- Model: MRI AGCM 3.1 (20 km-mesh)
- Projection periods:
  - Present-day exp. (PD) : 1979–2003 (25 yr)
  - Future global warmed exp. (GW): 2075–2099 (25 yr)
- Prescribed Lower Boundary Conditions:
  - PD: Observations (HadISST)
  - GW: Ensemble mean of 18 CMIP3 models' SST under the IPCC A1B scenario



- Relatively larger increase in SST in the Northern Hemisphere than in the Southern Hemisphere.
- The SST increase is the largest in the tropical Central Pacific.


# TC Detection Criteria

Based on Oouchi et al. (2006)


- Sea level pressure: **2.0 hPa** lower than the surroundings area.
- 850 hPa Relative vorticity:  **$3.0 \times 10^{-5} /s$**
- 850 hPa Maximum wind speed: **10.0 m/s**
- Warm Core: **1.0 K**
- Duration: **36 hours**
- Maximum wind speed at 850 hPa should be greater than the 300 hPa (to exclude extra-tropical cyclones).

# Simulated Global TC Tracks

Observations (1979-2003)




PD (1979-2003)



TC tracks are well simulated by the PD experiment.

# Future Changes in the Simulated Global TC Tracks

GW (2075-2099)




tern North Pacific).

(W-PD)


# What causes TC track changes?

## Steering flow (850-300hPa) changes



**Steering flow changes** (westerly flow anomaly) **partly** explain TC track changes by inhibiting westward TC motion.

## TC genesis frequency changes



**TC genesis location changes** (eastward shift) **mainly** explain TC track changes.



# Genesis potential index


To determine the factors behind such genesis changes, we used a Genesis Potential Index (GPI) by Emanuel and Nolan (2004) with some modifications.

$$GPI' = \left| 10^5 \eta \right|^{\frac{3}{2}} \left( \frac{RH}{50} \right)^3 \left( \frac{V_{pt}}{70} \right)^3 (1 + 0.1V_s)^{-2} \left( \frac{-\omega + 0.1}{0.1} \right),$$

Absolute Relative Maximum Vertical Wind Vertical Wind  
Vorticity Humidity Potential Shear (850- Velocity  
at 850hPa at 700hPa Intensity 200hPa) at 500hPa

Future changes in TC  
genesis frequency

GPI changes




Spatial correlation coefficient is 0.55.


GPI performs reasonably well in reflecting the changes in TC genesis frequency.

# Each term contribution to the changes in GPI

Relative humidity




Potential Intensity




**1. Thermodynamic changes has less influence.**

**=>Relative humidity and Potential intensity tend to cancel each other.**


Vorticity



Vertical Wind Shear




Vertical Wind Velocity



**2. Dynamical changes have great influences.**

**=>Vorticity and vertical wind shear contribute to the increase in GPI in the eastern WNP.**

**=>Vorticity and vertical wind velocity contribute to the decrease in GPI in the western WNP.**




# Mechanisms of future changes in TC genesis

## Future changes in vertical wind velocity at 500 hPa

### **Rossby wave response:**

Positive vorticity => Increase in TC genesis in the eastern WNP

Downward anomaly




### **Weakening of the Walker circulation:**

Decrease in upward motion => Decrease in TC genesis in the western WNP

Upward anomaly

# Why is upward motion enhanced in the tropical central Pacific?



Future changes in vertical motions (top) appear to be strongly related to the prescribed SST anomaly (bottom), indicating that spatial distribution of tropical SST may be a key factor for TC activities.


# Conclusion

The projected future changes in TC activity indicates:

- (a) Positions of the prevailing northward recurving **TC tracks will shift eastward** over the open ocean of the WNP;
- (b) **TC track changes are partially due to changes of the large scale steering flows**, but primarily **owing to the changes in TC-genesis locations**: TC formation will be less to the west of  $140^{\circ}\text{E}$ , whereas more in eastern WNP ( $0\text{--}17^{\circ}\text{N}$ ,  $140\text{--}180^{\circ}\text{E}$ );
- (c) **The decrease in the TC genesis frequency in the western WNP** are mainly due to *in situ* **reduction of the large scale ascent**, which is caused by the enhanced descending branch of the zonal circulation (i.e., weakening of the Walker circulation).
- (d) **The enhanced TC genesis in the eastern WNP** is due to the **increased in situ low-level cyclonic vorticity, reduced vertical wind shear**, caused by Rossby wave response induced by enhanced diabatic heating in the central tropical Pacific.

Thank you


# Future changes in “Landing” tropical cyclones




The frequency of tropical cyclones approaching Japan and Korea may decrease in the future.

However, once a TC approaches the coast lines, mean of maximum wind velocity is larger than the present climate, leading to a catastrophic damage in the future.

(a) July-October mean of TC frequency near coasts




(b) Mean maximum wind velocity of TCs near coasts






# Uncertainties caused by prescribed SSTs

## Individual SST Case 1




## Individual SST Case 2






# Robustness of WNP TC density changes among different experiments



# Interannual variations in TC number



Correlation Coefficients: PD and Obs = 0.55  
PD and GW = 0.20  
Obs and Nino3.4=0.27

# How to prescribe SST for the future experiment




FIG. Schematic diagram of prescribed lower boundary conditions for the future simulation.